

On the universality of the a priori mixing angles in weak radiative decays of hyperons

A. García

*Departamento de Física. Centro de Investigación y de Estudios Avanzados del IPN
A.P. 14-740. México, D.F. 07000. MEXICO*

R. Huerta

*Departamento de Física Aplicada
Centro de Investigación y de Estudios Avanzados del IPN, Unidad Mérida
A.P. 73, Cordemex. Mérida, Yucatán 97310. MEXICO*

G. Sánchez-Colón*

*Physics Department. University of California
Riverside, California 92521-0413. U.S.A.
(May 11, 1998)*

Abstract

Strong-flavor and parity a priori mixing in hadrons are shown to describe well the experimental evidence on weak radiative decays of hyperons. An independent determination of the a priori mixing angles is performed. The values obtained for them are seen to have a universality-like property, when compared to their values in non-leptonic decays of hyperons.

PACS number(s): 13.40.Hg, 11.30.Er, 11.30.Hr, 12.60.-i

Typeset using REVTeX

*Permanent Address: Departamento de Física Aplicada. Centro de Investigación y de Estudios Avanzados del IPN, Unidad Mérida. A.P. 73, Cordemex. Mérida, Yucatán 97310. MEXICO.

The enhancement phenomenon observed in weak decays of hadrons has remained an open challenge for quite sometime now. In a recent paper [1] we have studied the possibility of attributing this phenomenon to the existence of a priori strong-flavor and parity mixings in hadrons. In non-leptonic decays of hyperons (NLDH) we obtained two important results: (i) the predictions of the $|\Delta I| = 1/2$ rule and (ii) a fairly good description of the available experimental data. The a priori mixing angles, three in the case of NLDH which we named σ , δ , and δ' , appear as free parameters because we did not have any theoretical or phenomenological control on them. We obtained the values $\sigma = (4.9 \pm 1.5) \times 10^{-6}$, $|\delta| = (0.23 \pm 0.07) \times 10^{-6}$, and $|\delta'| = (0.26 \pm 0.07) \times 10^{-6}$. A more detailed analysis [2] indicates that the violations of the $|\Delta I| = 1/2$ rule affect their values and one should take more conservative ones as their estimates in NLDH, namely, $\sigma = (4.9 \pm 2.0) \times 10^{-6}$, $|\delta| = (0.22 \pm 0.09) \times 10^{-6}$, and $|\delta'| = (0.26 \pm 0.09) \times 10^{-6}$. The explicit absolute-values bars on δ and δ' are introduced as a reminder that their signs were not determined in NLDH.

The results obtained so far are very encouraging. Indeed, the a priori mixing scheme may serve as framework for the systematic description of the enhancement phenomenon. However, along this line, one faces very stringent tests that this scheme must pass successfully. Since the a priori mixings are non-perturbative effects in hadron wave functions, not only other decays processes originating from different dynamics where enhancement is present should be well described but the a priori angles σ , δ , and δ' should appear with values compatible with their values in NLDH, i.e., these angles must necessarily have a universality-like property. If this last is not obtained, then this should be sufficient reason to reject the whole scheme, at least as one with practical usefulness.

In this letter we shall make a detailed application of the above scheme to the weak radiative decays of hyperon (WRDH). The experimental evidence [3] on them is collected in column eight of Table I. If the enhancement phenomenon describes overwhelmingly these decays and if such phenomenon is described by a priori mixings, then these transition are due to the matrix elements of the ordinary parity and flavor conserving electromagnetic interaction hamiltonian $H_{em} = iJ_{em}^\mu A_\mu$ between a priori mixed baryons. The latter are given by

$$\begin{aligned}
p_{ph} &= p_{0s} - \sigma \Sigma_{0s}^+ - \delta \Sigma_{0p}^+ + \dots, \\
n_{ph} &= n_{0s} + \sigma \left(\frac{1}{\sqrt{2}} \Sigma_{0s}^0 + \sqrt{\frac{3}{2}} \Lambda_{0s} \right) + \delta \left(\frac{1}{\sqrt{2}} \Sigma_{0p}^0 + \sqrt{\frac{3}{2}} \Lambda_{0p} \right) + \dots, \\
\Sigma_{ph}^+ &= \Sigma_{0s}^+ + \sigma p_{0s} - \delta' p_{0p} + \dots, \\
\Sigma_{ph}^0 &= \Sigma_{0s}^0 + \frac{1}{\sqrt{2}} \sigma (\Xi_{0s}^0 - n_{0s}) + \frac{1}{\sqrt{2}} \delta \Xi_{0p}^0 + \frac{1}{\sqrt{2}} \delta' n_{0p} + \dots, \\
\Sigma_{ph}^- &= \Sigma_{0s}^- + \sigma \Xi_{0s}^- + \delta \Xi_{0p}^- + \dots, \\
\Lambda_{ph} &= \Lambda_{0s} + \sqrt{\frac{3}{2}} \sigma (\Xi_{0s}^0 - n_{0s}) + \sqrt{\frac{3}{2}} \delta \Xi_{0p}^0 + \sqrt{\frac{3}{2}} \delta' n_{0p} + \dots,
\end{aligned} \tag{1}$$

$$\Xi_{ph}^0 = \Xi_{0s}^0 - \sigma(\frac{1}{\sqrt{2}}\Sigma_{0s}^0 + \sqrt{\frac{3}{2}}\Lambda_{0s}) + \delta'(\frac{1}{\sqrt{2}}\Sigma_{0p}^0 + \sqrt{\frac{3}{2}}\Lambda_{0p}) + \dots,$$

$$\Xi_{ph}^- = \Xi_{0s}^- - \sigma\Sigma_{0s}^- + \delta'\Sigma_{0p}^- + \dots$$

In these expressions the subindices *naught*, *s*, and *p* indicate strong-flavor, positive, and negative parity eigenstates and each physical baryon is the mass eigenstate already observed [3]. A_μ represents the photon field operator and J_μ is the ordinary electromagnetic current operator. This operator is — and this must be emphasized — a flavor-conserving Lorentz proper four-vector. The hadronic parts of the transition amplitudes are then given by:

$$\langle p_{ph} | J_{em}^\mu | \Sigma_{ph}^+ \rangle = \bar{u}_p [\sigma(f_2^{\Sigma^+} - f_2^p) + (\delta' f_2^p - \delta f_2^{\Sigma^+}) \gamma^5] i\sigma^{\mu\nu} q_\nu u_{\Sigma^+}$$

$$\langle \Sigma_{ph}^- | J_{em}^\mu | \Xi_{ph}^- \rangle = \bar{u}_{\Sigma^-} [\sigma(f_2^{\Xi^-} - f_2^{\Sigma^-}) + (\delta' f_2^{\Sigma^-} - \delta f_2^{\Xi^-}) \gamma^5] i\sigma^{\mu\nu} q_\nu u_{\Xi^-}$$

$$\begin{aligned} \langle n_{ph} | J_{em}^\mu | \Lambda_{ph} \rangle = \bar{u}_n \left\{ \sigma \left[\sqrt{\frac{3}{2}}(f_2^\Lambda - f_2^n) + \frac{1}{\sqrt{2}}f_2^{\Sigma^0\Lambda} \right] \right. \\ \left. + \left[\sqrt{\frac{3}{2}}(\delta' f_2^n - \delta f_2^\Lambda) - \delta \frac{1}{\sqrt{2}}f_2^{\Sigma^0\Lambda} \right] \gamma^5 \right\} i\sigma^{\mu\nu} q_\nu u_\Lambda \end{aligned}$$

$$\begin{aligned} \langle \Lambda_{ph} | J_{em}^\mu | \Xi_{ph}^0 \rangle = \bar{u}_\Lambda \left\{ \sigma \left[\sqrt{\frac{3}{2}}(f_2^{\Xi^0} - f_2^\Lambda) - \frac{1}{\sqrt{2}}f_2^{\Sigma^0\Lambda} \right] \right. \\ \left. + \left[\sqrt{\frac{3}{2}}(\delta' f_2^\Lambda - \delta f_2^{\Xi^0}) + \delta' \frac{1}{\sqrt{2}}f_2^{\Sigma^0\Lambda} \right] \gamma^5 \right\} i\sigma^{\mu\nu} q_\nu u_{\Xi^0} \end{aligned}$$

$$\begin{aligned} \langle \Sigma_{ph}^0 | J_{em}^\mu | \Xi_{ph}^0 \rangle = \bar{u}_{\Sigma^0} \left\{ \sigma \left[\frac{1}{\sqrt{2}}(f_2^{\Xi^0} - f_2^{\Sigma^0}) - \sqrt{\frac{3}{2}}f_2^{\Sigma^0\Lambda} \right] \right. \\ \left. + \left[\frac{1}{\sqrt{2}}(\delta' f_2^{\Sigma^0} - \delta f_2^{\Xi^0}) + \delta' \sqrt{\frac{3}{2}}f_2^{\Sigma^0\Lambda} \right] \gamma^5 \right\} i\sigma^{\mu\nu} q_\nu u_{\Xi^0} \end{aligned} \quad (2)$$

In these amplitudes only contributions to first order in σ , δ , and δ' need be kept. Each matrix element is flavor and parity conserving and can be expanded in terms of charge $f_1(0)$ form factors and anomalous magnetic $f_2(0)$ form factors. Because the charges of the positive- and negative-parity parts of the same physical wave function are equal and such changes are controlled by the generator property of J_μ all the f_1 's cancel away and only the f_2 contribute. The f_2 between *s* and *p* parts and between *p* and *s* parts can be identified with the f_2 between *s* and *s* parts, provided that a relative minus sign be present between the former two in order to respect hermicity. Notice that the amplitudes (2) are all of the form $\bar{u}_B(C + D\gamma_5)i\sigma^{\mu\nu}q_\nu u_A$, where C is the so-called parity conserving amplitude and D is

the so-called parity violating one. We stress, however, that in this model both C and D are parity and flavor conserving.

In principle, we have information about all the quantities that appear in these amplitudes for WRDH. The a priori angles are known from NLDH and the f_2 can be related to the measured total magnetic moments of spin 1/2 baryons. The latter values are displayed in the second column of Table I [3]. However, it is important that the mixing angles be determined independently in WRDH and, accordingly, we shall use them as free parameters in the remaining of this paper. How the f_2 's are related to the observed total magnetic moments is a question we shall deal with in steps. As a first approximation we shall assume that the f_2 's are related to the μ 's of Table I by the formula $\mu_A^{exp} = e_A + f_2^A$ where A is a baryon and e_A its charge. Thus, for example, f_2^p obeys the relationship $\mu_p^{exp} = 1 + f_2^p$ (in nuclear magnetons), etc. Using this assumption we may compare with the experimental data of WRDH. The predictions obtained are displayed in the columns I of Table I.

These first results are not quite good yet but they have a qualitative value. The main point is that the a priori mixing angles come out with the same order of magnitude observed in NLDH, which is very encouraging. The predictions for the observables are some very good, some good, but some show important deviations. The latter still have qualitative value, but should be improved. The values of the μ 's agree fairly well with their experimental counterparts.

As an intermediate step in this analysis it turns out to be very helpful to see what are the values of the total magnetic moments required to reproduce well the experimental observables of WRDH. This is achieved by relaxing the error bars of the measured μ 's up to 10% of the corresponding central values and, then, repeating the previous step. The results are displayed in the columns II of Table I. The experimental data are very well reproduced now, but at the expense of sizable (several percent) changes in the μ 's and new values for the mixing angles.

This second step clearly shows that we must accept that our first approximation—that of identifying the experimentally measured μ 's with the ones that are actually related to the f_2 's in this approach to WRDH—must be improved. The μ 's to be used for determining the f_2 's in the WRDH amplitudes are really transition magnetic moments. For example, the measured value of μ_p corresponds to the matrix element $\langle p_{ph} | J_{em}^\mu | p_{hp} \rangle \simeq \langle p_{os} | J_{em}^\mu | p_{os} \rangle$, where both physical wave functions carry the mass m_p . In contrast, the μ_p that appears in $\Sigma^+ \rightarrow p\gamma$ corresponds to a matrix element whose bra carries the mass m_p and whose ket carries the mass m_{Σ^+} . So, the normalization of μ_p originating in the matrix element $\langle p_{ph} | J_\mu | \Sigma_{ph}^+ \rangle$ should be related to both masses, m_p and m_{Σ^+} . It is in this sense that the magnetic moments that we must use are transition magnetic moments.

The natural normalization of magnetic moments is determined by the Gordon decomposition [4]. Using this expansion for guidance, then, for example, μ_p should be normalized to $m_p + m_{\Sigma^+}$ and not to $2m_p$, etc. One can see already a qualitative indication of this happening in the first column II in Table I, the changes in the μ 's are systematically in this direction. μ_p , μ_n , μ_{Ξ^-} , μ_{Σ^-} , μ_{Σ^+} , μ_{Ξ^0} , and μ_{Σ^0} appear to become smaller or larger according to such changes in normalization. μ_Λ and $\mu_{\Sigma^0\Lambda}$ are mixed cases because they appear in two or three decays and will be required to be reduced or to be increased in going from one case to another and, therefore, Table I cannot provide a clear cut tendency.

Our third step is to improve our approximation following the above discussion. One must

change the normalization of the total magnetic moments either by applying, for example, the factor $(m_p + m_p)/(m_p + m_{\Sigma^+})$ to the experimental μ_p or the inverse factor to the theoretical μ_p related to f_2^p . Numerically, either way leads to the same result. For definiteness, we choose the former. The corrected experimental values are displayed in column five of Table I. Then, recalculating everything lead to the predictions of columns labeled III of Table I. The values of the mixing angles appear in the bottom of the last column of this table.

The overall agreement is greatly improved, the experimental data are well produced while keeping the magnetic moments in very good agreement with their experimental counterparts. The only deviations that merit further discussion appear in Γ_2 and in μ_{Σ^-} which are intimately related. These deviations are probably due to another one of our approximations: ignoring the contributions of W_μ [5]. We have assumed these contributions to be small (non-enhanced) and we have neglected them, but since they are non-zero they should eventually be included. Their effect should appear mainly in small amplitudes. Γ_2 is about an order of magnitude smaller than other Γ 's and, accordingly, its corresponding amplitudes are also the smallest ones. The interference of the W_μ amplitudes should be relatively more important in this case. Therefore, one is not entitled to expect a better agreement of a priori mixings in this case. The agreement already obtained is probably the best one can hope for if one stays short of actually incorporating the contributions of W_μ , as we have.

In Table I we included the predictions for α_2 and α_3 . They are also quite sensible to the a priori mixing parameters. It would be important to have experimental information on them. At present only $\alpha_2 = 1.0 \pm 1.3$ is available [6]. This measurement can be understood to mean that only its sign is known, namely, positive. The prediction in case III agrees with this sign.

We are now in a position to conclude our present analysis. To extend the credibility of the a priori mixing scheme it was very important to be able to describe WRDH. As we have shown above this is achieved. However, the most important and stringent test is that the mixing angles share a universality-like property. The values for them obtained independently in WRDH are in very good agreement with the absolute values obtained for them in NLDH. It is the passing this universality-like test that lends the strongest support to the possibility that the above scheme may serve a framework for the systematic description of the enhancement phenomenon in weak decays of hadrons.

As a closing remark, let us mention how the three orders of magnitudes observed experimentally between NLDH and WRDH can be understood: The enhancement phenomenon dominates these decays, such phenomenon is described by the a priori mixings scheme, and the mixing angles are common to both dynamics, then, the relative order of magnitude between these two sets of decays is given by the ratio $(g^2/4\pi)/(e^2/4\pi) \sim 10^3$, where g is the strong-interaction Yukawa coupling constant [7].

The authors wish to acknowledge partial support by CONACyT (México).

REFERENCES

- [1] A. García, R. Huerta, and G. Sánchez-Colón, “*A priori* mixed hadrons, weak radiative and non-leptonic decays of hyperons”. e-print archive hep-ph/9804341, to be published in J. of Phys. **G**.
- [2] A. García, R. Huerta, and G. Sánchez-Colón, “A detailed determination of the *a priori* mixing angles in non-leptonic decays of hyperons”. e-print archive hep-ph/9804340, submitted to J. of Phys. **G**.
- [3] For a review of particle properties, see L. Montanet *et al.* Phys. Rev. **D 50**, 1173 (1994).
- [4] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Mechanics* (McGraw-Hill, New York, 1964), Chap. 3.
- [5] A recent reviews of the W_μ contributions to WRDH are found in B. Bassalleck, Nucl. Phys. A **547**, 299c (1992) and in J. Lach and P. Żenczykowski, Int. J. Mod. Phys. **A 10**, 3817 (1995).
- [6] See J. Lach and P. Żenczykowski in Ref. [5].
- [7] O. Dumbrajs *et al.* Nucl. Phys. **B 216**, 277 (1983).

TABLES

TABLE I. Experimental values of the observables in WRDH and of the magnetic moments (m. m.) of hyperons along with the predictions of the three cases considered. The indices 1,2,...,5 on the observables correspond, respectively, to the decays $\Sigma^+ \rightarrow p\gamma$, $\Xi^- \rightarrow \Sigma^-\gamma$, $\Lambda \rightarrow n\gamma$, $\Xi^0 \rightarrow \Lambda\gamma$, and $\Xi^0 \rightarrow \Sigma^0\gamma$. The numbers in parenthesis in the m. m. indicate the decay in which they appear. The values of the a priori mixing angles in column eight come from NLDH. The mixing angles are in 10^{-6} , the decay rates are in 10^6 sec^{-1} , and the m. m. are in nuclear magnetons. The only m. m. that has not been measured is μ_{Σ^0} . We have taken for it its $SU(3)$ estimate with a 10% error bar.

	Magnetic moments			Transition m. m.			Observables and angles			
	Exp.	I	II	Exp.	III		Exp.	I	II	III
$\mu_p(1)$	2.793	2.793	2.745	2.463	2.463	Γ_1	15.65 ± 0.88	11.55	15.60	14.62
$\mu_n(3)$	-1.913	-1.913	-1.654	-1.750	-1.750	Γ_2	0.77 ± 0.14	1.37	0.81	1.30
$\mu_{\Xi^-}(2)$	-0.651 ± 0.003	-0.652	-0.747	-0.683 ± 0.003	-0.685	Γ_3	6.65 ± 0.57	7.13	6.67	6.16
$\mu_{\Sigma^-}(2)$	-1.160 ± 0.025	-1.018	-0.868	-1.103 ± 0.024	-0.958	Γ_4	3.66 ± 0.56	5.26	3.68	4.38
$\mu_{\Lambda}(3)$	-0.613 ± 0.004	-0.611	-0.553	-0.665 ± 0.004	-0.665	Γ_5	12.1 ± 1.4	5.23	11.8	10.13
$\mu'_{\Lambda}(4)$				-0.563 ± 0.004	-0.562	α_1	-0.76 ± 0.08	-0.78	-0.75	-0.88
$\mu_{\Sigma^+}(1)$	2.458 ± 0.010	2.430	2.553	2.748 ± 0.011	2.763	α_2		-0.09	0.56	0.20
$\mu_{\Xi^0}(4)$	-1.250 ± 0.014	-1.271	-1.502	-1.353 ± 0.015	-1.357	α_3		-0.79	-0.83	0.76
$\mu'_{\Xi^0}(5)$				-1.311 ± 0.015	-1.305	α_4	0.40 ± 0.40	-0.86	0.18	0.27
$\mu_{\Sigma^0\Lambda}(3)$	-1.610 ± 0.080	-1.597	-1.626	-1.808 ± 0.090	-1.659	α_5	0.20 ± 0.32	-0.46	-0.02	0.77
$\mu'_{\Sigma^0\Lambda}(4)$				-1.529 ± 0.076	-1.426	σ	4.9 ± 2.0	0.70 ± 0.03	1.72 ± 0.46	0.91 ± 0.06
$\mu''_{\Sigma^0\Lambda}(5)$				-1.482 ± 0.074	-1.617	δ	$ 0.22 \pm 0.09$	0.03 ± 0.08	-0.40 ± 0.19	-0.11 ± 0.08
$\mu_{\Sigma^0}(5)$	0.649 ± 0.065	0.499	0.624	0.617 ± 0.062	0.500	δ'	$ 0.26 \pm 0.09$	0.10 ± 0.07	-0.28 ± 0.18	-0.25 ± 0.08